

HUTT ESTUARY: INTERTIDAL SEDIMENT MONITORING SUMMARY, 2013/2014

Prepared for Greater Wellington Regional Council by Leigh Stevens and Barry Robertson, Wriggle Coastal Management, March 2014



Figure 1. Location of intertidal sediment plates and fine scale monitoring sites in the lower Hutt Estuary.








Figure 3. Sediment mud content (+/-SE, n=3), Hutt Estuary, 2010-12, 2014^{*}. *2014 = single composite sample

This summary card presents the results of monitoring undertaken on 22 January 2014 to track changes to sediment indicators in Hutt Estuary. Detailed reporting is scheduled to be undertaken 5 yearly (next due 2017).

Methods

The depths to four concrete plates buried in intertidal sediment in 2010 were measured to assess the long-term sedimentation rate (Figure 1 - see Robertson and Stevens 2011 for full details). Redox Potential Discontinuity (RPD) depth and sediment grain size were assessed to indicate sediment condition.

Risk Indicator Ratings

To help quickly identify the potential significance of sediment to Hutt Estuary, "risk indicator ratings" have been proposed (Table 1, see Stevens and Robertson 2014 for further detail) and are part of a suite of indicators being developed to assess the predominant issues affecting NZ estuaries (i.e. eutrophication, sedimentation, disease risk, toxicity and habitat change -Robertson and Stevens 2006, 2012, 2013). For each indicator, relative levels of risk (e.g. very low, low, moderate, high, very high) are assigned based on their relationship with water or sediment quality. Each rating is designed to be used in combination with relevant information and other risk indicator ratings, and under expert guidance, to assess overall estuary condition in relation to key issues, and monitoring and management recommendations.

Table 1. Risk indicator ratings for sedimentation rate, sediment mud content, and RPD depth.

RISK INDICATOR RATING	SEDIMENTATION RATE	MUD CONTENT*	RPD DEPTH	
Very Low	<1mm/yr	<2%	>10cm	
Low	>1-2mm/yr	2-5%	3-10cm	
Moderate	>2-5mm/yr	>5-15%	1-<3cm	
High	>5-10mm/yr	>15-25%	0-<1cm	
Very High	>10mm/yr	>25%	Anoxic at surface	

* rating revised in 2014 based on Robertson (2013).

2010-2014 Sedimentation Rate

Figure 2 and Table 2 summarise sediment level changes since the 2010 baseline. Changes in sediment levels over individual plates range from -13 to +2mm/yr, with an annual site average range of -0.8 to -9.3mm/yr. The overall mean sedimentation rate across the four years of monitoring is a decrease of 4.2mm/yr. Regular dredging of sediments from the channel in the lower estuary, and scouring of tidal flats during high river flows, are likely reasons for the low mean annual deposition rate recorded at the monitoring site.

2014 Sediment Mud Content and RPD depth

Sediment mud content was 21.9% (Table 3), reflecting firm muddy sands. The mud content is at the lower end of the 23-51% range recorded from the nearby shallow subtidal fine scale sites (Figures 1 and 3). Average RPD depth was 1.5cm (Table 3). RPD and mud content fall within the "moderate" and "high" risk indicator ratings respectively.

Conclusion

The sedimentation rate over the past 4 years showed slight erosion, but the high sediment mud content and shallow RPD depth indicate the estuary is susceptible to sediment related impacts from poor clarity and muddy intertidal substrates, with a macrofaunal community dominated by mud tolerant species - a common situation in NZ tidal river estuaries.

Recommended Monitoring

Continue annual monitoring of sediment rate, RPD and grain size to measure sediment deposition and temporal change on the only significant remaining intertidal flat within the estuary. Report results annually via a summary card, with detailed reporting undertaken 5 yearly in conjunction with fine scale monitoring (next scheduled for 2017).

HUTT ESTUARY: INTERTIDAL SEDIMENT MONITORING SUMMARY, 2013/2014

CITE	Measured Mean Depth to Sediment Plate (mm)			Change in Sediment Level Over Plate (mm)			SEDIMENTATION RATE 2010-2014				
SITE	11 Apr 2010	15 Jan 2011	21 Feb 2012	15 Jan 2013	22/01/2014	2010-2011	2011-2012	2012-2013	2013-2014	(mm/yr)	RISK RATING
Hutt Plate 1	257	256	247	246	240	-1	-9	-1	-6	-4.2 (SE=0.33)	VERY LOW
Hutt Plate 2	250	248	245	242	232	-2	-3	-3	-10		
Hutt Plate 3	295	297	290	289	276	+2	-7	-1	-13		
Hutt Plate 4	287	285	285	282	274	-2	0	-3	-8		
	Mean Change in Sediment Level (mm/yr)			-0.8	-4.8	-2.0	-9.3				

Table 2. Sediment monitoring results for Hutt Estuary, April 2010 - January 2014.

Table 3. Grain size results for the Hutt Estuarysedimentation plate site, 22 January 2014.

Date	RPD mean depth	Mud	Sand	Gravel
22 Jan 2014	1.5 (SE=0.2)	21.9%	74.5%	3.6%

Note: Grain size results are based on a single composite sample comprising 10 sub-samples collected from the site. Mean RPD depth is derived from 10 replicate measures.

References

- Robertson, B.P. 2013. Determining the sensitivity of macroinvertebrates to fine sediments in representative New Zealand estuaries. Honours thesis, University of Victoria, Wellington.
- Robertson, B.M. and Stevens, L. 2006. Southland Estuaries State of Environment Report 2001-2006. Prepared for Environment Southland. 45p plus appendices.
- Robertson, B.M. and Stevens, L. 2011. Hutt Estuary: Fine Scale Monitoring 2010/11. Prepared for Greater Wellington Regional Council. 25p.
- Robertson, B.M. and Stevens, L. 2012. Tasman Coast: Waimea Inlet to Kahurangi Point, habitat mapping, risk assessment and monitoring recommendations. Prepared for Tasman District Council. 167p.
- Robertson, B.M. and Stevens, L. 2013. Moutere Inlet fine scale monitoring 2012/2013. Prepared for Tasman District Council. 25p.
- Stevens, L. and Robertson, B.M. 2014. Whareama Estuary: Intertidal Sediment Monitoring 2013/14. Prepared for Greater Wellington Regional Council. 6p.

Location of sedimentation rate monitoring plates in Hutt Estuary.

Site	NZTM East	NZTM North
Plate 1	1759101	5433548
Plate 2	1759097	5433548
Plate 3	1759093	5433548
Plate 4	1759089	5433548
Peg 1	1759103	5433548
Peg 2	1759099	5433548
Peg 3	1759095	5433548
Peg 4	1759091	5433548
Peg 5	1759087	5433548



Intertidal flats in the lower Hutt Estuary, Jan 2014